

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTAMXG1614

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	FEB 28	PATDPAFULL - New display fields provide for legal status data from INPADO
NEWS	4	FEB 28	BABS - Current-awareness alerts (SDIs) available
NEWS	5	MAR 02	GBFULL: New full-text patent database on STN
NEWS	6	MAR 03	REGISTRY/ZREGISTRY - Sequence annotations enhanced
NEWS	7	MAR 03	MEDLINE file segment of TOXCENTER reloaded
NEWS	8	MAR 22	KOREAPAT now updated monthly; patent information enhanced
NEWS	9	MAR 22	Original IDE display format returns to REGISTRY/ZREGISTRY
NEWS	10	MAR 22	PATDPASPC - New patent database available
NEWS	11	MAR 22	REGISTRY/ZREGISTRY enhanced with experimental property tags
NEWS	12	APR 04	EPFULL enhanced with additional patent information and new fields
NEWS	13	APR 04	EMBASE - Database reloaded and enhanced
NEWS	14	APR 18	New CAS Information Use Policies available online
NEWS	15	APR 25	Patent searching, including current-awareness alerts (SDIs), based on application date in CA/CAPLUS and USPATFULL/USPAT2 may be affected by a change in filing date for U.S. applications.
NEWS	16	APR 28	Improved searching of U.S. Patent Classifications for U.S. patent records in CA/CAPLUS
NEWS	17	MAY 23	GBFULL enhanced with patent drawing images
NEWS	18	MAY 23	REGISTRY has been enhanced with source information from CHEMCATS
NEWS	19	JUN 06	The Analysis Edition of STN Express with Discover! (Version 8.0 for Windows) now available
NEWS	20	JUN 13	RUSSIAPAT: New full-text patent database on STN
NEWS	21	JUN 13	FRFULL enhanced with patent drawing images
NEWS	22	JUN 27	MARPAT displays enhanced with expanded G-group definitions and text labels
NEWS	23	JUL 01	MEDICONF removed from STN
NEWS	24	JUL 07	STN Patent Forums to be held in July 2005
NEWS	25	JUL 13	SCISEARCH reloaded
NEWS	26	JUL 20	Powerful new interactive analysis and visualization software, STN AnaVist, now available
NEWS	27	AUG 11	Derwent World Patents Index(R) web-based training during August
NEWS	28	AUG 11	STN AnaVist workshops to be held in North America
NEWS	EXPRESS		JUNE 13 CURRENT WINDOWS VERSION IS V8.0, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 13 JUNE 2005
NEWS	HOURS		STN Operating Hours Plus Help Desk Availability
NEWS	INTER		General Internet Information
NEWS	LOGIN		Welcome Banner and News Items
NEWS	PHONE		Direct Dial and Telecommunication Network Access to STN
NEWS	WWW		CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 13:19:44 ON 22 AUG 2005

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 13:19:54 ON 22 AUG 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 19 AUG 2005 HIGHEST RN 861198-35-8

DICTIONARY FILE UPDATES: 19 AUG 2005 HIGHEST RN 861198-35-8

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when conducting SmartSELECT searches.

*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*



Structure search iteration limits have been increased. See HELP SLIMITS for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:

<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Documents and Settings\mgraffeo\My Documents\Critical Data\10517801\compound.str



chain nodes :
1 2 3 4 5 6 7 8 9 10 11 12 13 14
chain bonds :
1-2 2-3 2-12 3-4 4-5 5-6 6-7 7-8 8-9 9-10 9-14 10-11 10-13
exact/norm bonds :
2-3 2-12 3-4 5-6 6-7 9-14
exact bonds :
1-2 4-5 7-8 8-9 9-10
normalized bonds :
10-11 10-13

Match level :
1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS
10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS

Stereo Bonds:

14-9 (Single Hash).

Stereo Chiral Centers:

9 (Parity=Don't Care)

Stereo RSS Sets:

Type=Relative (Default). 1 Nodes= 9

L1 STRUCTURE UPLOADED

=> s l1 sss full
FULL SEARCH INITIATED 13:20:10 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 131 TO ITERATE

100.0% PROCESSED 131 ITERATIONS 16 ANSWERS
SEARCH TIME: 00.00.01

L2 16 SEA SSS FUL L1

=> s l1 exa full
FULL SEARCH INITIATED 13:20:20 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 8 TO ITERATE

100.0% PROCESSED 8 ITERATIONS 2 ANSWERS
SEARCH TIME: 00.00.01

L3 2 SEA EXA FUL L1

=> file caplus
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY SESSION
FULL ESTIMATED COST 215.58 215.79

FILE 'CAPLUS' ENTERED AT 13:20:28 ON 22 AUG 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 22 Aug 2005 VOL 143 ISS 9
FILE LAST UPDATED: 21 Aug 2005 (20050821/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l2 or l3

16 L2

13 L3

L4 16 L2 OR L3

=> s l4 (L) antioxidant

99349 ANTIOXIDANT

L5 0 L4 (L) ANTIOXIDANT

=> s l4 and chelat?

126712 CHELAT?

L6 1 L4 AND CHELAT?

=> d bib abs hitstr

L6 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2005 ACS on STN

AN 2004:2680 CAPLUS

DN 140:65201

TI (2S)-2-Amino-4-{[2-(ethanimidoylamino)ethyl]thio}butanoic acid nitric oxide synthase inhibitor in stabilized pharmaceutical dosage forms

IN Broughton, Stuart James; Gharu, Rajinder Kumar; Leow, Mark Yuen Tuck; Neale, Philip John

PA SB Pharmco Puerto Rico Inc., P. R.

SO PCT Int. Appl., 16 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004000296	A1	20031231	WO 2003-EP6465	20030619
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	EP 1513511	A1	20050316	EP 2003-740281	20030619
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			

PRAI GB 2002-14147 A 20020619

WO 2003-EP6465 W 20030619

AB Pharmaceutical compns. comprising (2S)-MeC(:NH)NHCH₂CH₂SCH₂CH₂CH(NH₂)CO₂H (I) a pharmaceutically acceptable bulking agent and one or more antioxidants or chelating agents are described. A direct compression formula for tablets contained I, EDTA, Avical PH101, silica, and Mg stearate.


IT 210354-22-6 438542-15-5 638198-40-0
638198-41-1

RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
((2S)-2-Amino-4-{ [2-(ethanimidoethylamino)ethyl]thio}butanoic acid nitric
oxide synthase inhibitor in stabilized pharmaceutical dosage forms)

RN 210354-22-6 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 438542-15-5 CAPLUS


CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]-, phosphate (1:1) (9CI)
(CA INDEX NAME)

CM 1

CRN 210354-22-6

CMF C8 H17 N3 O2 S


Absolute stereochemistry.



CM 2

CRN 7664-38-2


CMF H3 O4 P



RN 638198-40-0 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]-, monohydrate (9CI) (CA
INDEX NAME)


Absolute stereochemistry.



● H₂O

RN 638198-41-1 CAPLUS
 CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]-, trihydrate (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● 3 H₂O

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s 14 (L) (edta or "ethylenediamintetraacetic acid" or malic or ascorbic)
 82261 EDTA
 12 "ETHYLENEDIAMINTETRAACETIC"
 4020182 "ACID"
 11 "ETHYLENEDIAMINTETRAACETIC ACID"
 ("ETHYLENEDIAMINTETRAACETIC" (W) "ACID")
 29892 MALIC
 78739 ASCORBIC
 L7 0 L4 (L) (EDTA OR "ETHYLENEDIAMINTETRAACETIC ACID" OR MALIC OR
 ASCORBIC)

=>

---Logging off of STN---

=>
 Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	20.42	236.21
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL

	ENTRY	SESSION
CA SUBSCRIBER PRICE	-0.73	-0.73

STN INTERNATIONAL LOGOFF AT 13:23:20 ON 22 AUG 2005

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTAMXG1614

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1			Web Page URLs for STN Seminar Schedule - N. America
NEWS	2			"Ask CAS" for self-help around the clock
NEWS	3	FEB	28	PATDPAFULL - New display fields provide for legal status data from INPADOC
NEWS	4	FEB	28	BABS - Current-awareness alerts (SDIs) available
NEWS	5	MAR	02	GBFULL: New full-text patent database on STN
NEWS	6	MAR	03	REGISTRY/ZREGISTRY - Sequence annotations enhanced
NEWS	7	MAR	03	MEDLINE file segment of TOXCENTER reloaded
NEWS	8	MAR	22	KOREAPAT now updated monthly; patent information enhanced
NEWS	9	MAR	22	Original IDE display format returns to REGISTRY/ZREGISTRY
NEWS	10	MAR	22	PATDPASPC - New patent database available
NEWS	11	MAR	22	REGISTRY/ZREGISTRY enhanced with experimental property tags
NEWS	12	APR	04	EPFULL enhanced with additional patent information and new fields
NEWS	13	APR	04	EMBASE - Database reloaded and enhanced
NEWS	14	APR	18	New CAS Information Use Policies available online
NEWS	15	APR	25	Patent searching, including current-awareness alerts (SDIs), based on application date in CA/Caplus and USPATFULL/USPAT2 may be affected by a change in filing date for U.S. applications.
NEWS	16	APR	28	Improved searching of U.S. Patent Classifications for U.S. patent records in CA/Caplus
NEWS	17	MAY	23	GBFULL enhanced with patent drawing images
NEWS	18	MAY	23	REGISTRY has been enhanced with source information from CHEMCATS
NEWS	19	JUN	06	The Analysis Edition of STN Express with Discover! (Version 8.0 for Windows) now available
NEWS	20	JUN	13	RUSSIAPAT: New full-text patent database on STN
NEWS	21	JUN	13	FRFULL enhanced with patent drawing images
NEWS	22	JUN	27	MARPAT displays enhanced with expanded G-group definitions and text labels
NEWS	23	JUL	01	MEDICONF removed from STN
NEWS	24	JUL	07	STN Patent Forums to be held in July 2005
NEWS	25	JUL	13	SCISEARCH reloaded
NEWS	26	JUL	20	Powerful new interactive analysis and visualization software, STN AnaVist, now available
NEWS	27	AUG	11	Derwent World Patents Index(R) web-based training during August
NEWS	28	AUG	11	STN AnaVist workshops to be held in North America
NEWS EXPRESS				JUNE 13 CURRENT WINDOWS VERSION IS V8.0, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 13 JUNE 2005

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 14:57:13 ON 22 AUG 2005

=> file reg

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 14:57:25 ON 22 AUG 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 19 AUG 2005 HIGHEST RN 861198-35-8

DICTIONARY FILE UPDATES: 19 AUG 2005 HIGHEST RN 861198-35-8

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when conducting SmartSELECT searches.



*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*

Structure search iteration limits have been increased. See HELP SLIMITS for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Documents and Settings\mgraffeo\My Documents\Critical Data\10517801\compound.str



chain nodes :

1 2 3 4 5 6 7 8 9 10 11 12 13 14

chain bonds :

1-2 2-3 2-12 3-4 4-5 5-6 6-7 7-8 8-9 9-10 9-14 10-11 10-13

exact/norm bonds :

2-3 2-12 3-4 5-6 6-7 9-14

exact bonds :

1-2 4-5 7-8 8-9 9-10

normalized bonds :

10-11 10-13

Match level :

1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS
10:CLASS 11:CLASS 12:CLASS 13:CLASS 14:CLASS

Stereo Bonds:

14-9 (Single Hash).

Stereo Chiral Centers:

9 (Parity=Don't Care)

Stereo RSS Sets:

Type=Relative (Default). 1 Nodes= 9

L1 STRUCTURE UPLOADED

=> s l1 exa full

FULL SEARCH INITIATED 14:57:50 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 8 TO ITERATE

100.0% PROCESSED 8 ITERATIONS

2 ANSWERS

SEARCH TIME: 00.00.01

L2 2 SEA EXA FUL L1

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

54.68

54.89

FILE 'CAPLUS' ENTERED AT 14:57:58 ON 22 AUG 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 22 Aug 2005 VOL 143 ISS 9
FILE LAST UPDATED: 21 Aug 2005 (20050821/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l2 and (malic or edta or edetic or ascorbic)

13 L2
29892 MALIC
82261 EDTA
227 EDETIC
78739 ASCORBIC

L3 1 L2 AND (MALIC OR EDTA OR EDETIC OR ASCORBIC)


=> d bib abs hitstr

L3 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2005 ACS on STN
AN 2004:2680 CAPLUS
DN 140:65201
TI (2S)-2-Amino-4-{[2-(ethanimidoylamino)ethyl]thio}butanoic acid nitric
oxide synthase inhibitor in stabilized pharmaceutical dosage forms
IN Broughton, Stuart James; Gharu, Rajinder Kumar; Leow, Mark Yuon Tuck;
Neale, Philip John
PA SB Pharmco Puerto Rico Inc., P. R.
SO PCT Int. Appl., 16 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004000296	A1	20031231	WO 2003-EP6465	20030619
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
	EP 1513511	A1	20050316	EP 2003-740281	20030619
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
PRAI	GB 2002-14147	A	20020619		
	WO 2003-EP6465	W	20030619		
AB	Pharmaceutical compns. comprising (2S)-MeC(:NH)NHCH ₂ CH ₂ SCH ₂ CH ₂ CH(NH ₂)CO ₂ H (I) a pharmaceutically acceptable bulking agent and one or more antioxidants or chelating agents are described. A direct compression formula for tablets contained I, EDTA, Avical PH101, silica, and Mg stearate.				
IT	210354-22-6	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses) (2S)-2-Amino-4-{[2-(ethanimidoylamino)ethyl]thio}butanoic acid nitric oxide synthase inhibitor in stabilized pharmaceutical dosage forms)			
RN	210354-22-6	CAPLUS			

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s l2 and stability

13 L2

623742 STABILITY

L4 0 L2 AND STABILITY

=> s l2 and formulation

13 L2

126374 FORMULATION

L5 0 L2 AND FORMULATION

=> s l2 (antioxidant or chelat?)

MISSING OPERATOR 'L2 (ANTIOXIDAN'

The search profile that was entered contains terms or
nested terms that are not separated by a logical operator.

=> s l2 and (antioxidant or chelat?)

13 L2

99349 ANTIOXIDANT

126712 CHELAT?

L6 1 L2 AND (ANTIOXIDANT OR CHELAT?)

=> d bib abs

L6 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2005 ACS on STN

AN 2004:2680 CAPLUS

DN 140:65201

TI (2S)-2-Amino-4-{[2-(ethanimidoylamino)ethyl]thio}butanoic acid nitric
oxide synthase inhibitor in stabilized pharmaceutical dosage forms

IN Broughton, Stuart James; Gharu, Rajinder Kumar; Leow, Mark Yuen Tuck;
Neale, Philip John

PA SB Pharmco Puerto Rico Inc., P. R.

SO PCT Int. Appl., 16 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
	-----	---	-----	-----	-----
PI	WO 2004000296	A1	20031231	WO 2003-EP6465	20030619
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			

EP 1513511 A1 20050316 EP 2003-740281 20030619
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK

PRAI GB 2002-14147 A 20020619
WO 2003-EP6465 W 20030619

AB Pharmaceutical compns. comprising (2S)-MeC(:NH)NHCH₂CH₂SCH₂CH₂CH(NH₂)CO₂H
(I) a pharmaceutically acceptable bulking agent and one or more
antioxidants or chelating agents are described. A direct
compression formula for tablets contained I, EDTA, Avical PH101, silica,
and Mg stearate.

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s 12

L7 13 L2

=> d 1-13 bib abs

L7 ANSWER 1 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN

AN 2005:465475 CAPLUS

DN 143:71325

TI GW274150 and GW273629 are potent and highly selective inhibitors of
inducible nitric oxide synthase in vitro and in vivo

AU Alderton, Wendy K.; Angell, Anthony D. R.; Craig, Caroline; Dawson, John;
Garvey, Edward; Moncada, Salvador; Monkhouse, Jayne; Rees, Daryl; Russell,
Linda J.; Russell, Rachel J.; Schwartz, Sheila; Waslidge, Neil; Knowles,
Richard G.

CS Medicines Research Centre, Respiratory & Inflammation Centre of Excellence
for Drug Discovery, GlaxoSmithKline Research, Stevenage, SG1 2NY, UK

SO British Journal of Pharmacology (2005), 145(3), 301-312

CODEN: BJPCBM; ISSN: 0007-1188

PB Nature Publishing Group

DT Journal

LA English

AB GW274150 ([2-[(1-iminoethyl)amino]ethyl]-L-homocysteine) and GW273629
(3-[[2-[(1-iminoethyl)amino]ethyl]sulfonyl]-L-alanine) are potent,
time-dependent, highly selective inhibitors of human inducible nitric
oxide synthase (iNOS) vs. endothelial NOS (eNOS) (>100-fold) or neuronal
NOS (nNOS) (>80-fold). GW274150 and GW273629 are arginine competitive,
NADPH-dependent inhibitors of human iNOS with steady state K_d values of
<40 and <90 nM, resp. GW274150 and GW273629 inhibited intracellular iNOS
in J774 cells in a time-dependent manner, reaching IC₅₀ values of
0.2±0.04 and 1.3±0.16 µM, resp. They were also acutely selective
in intact rat tissues: GW274150 was >260-fold and 219-fold selective for
iNOS against eNOS and nNOS, resp., while GW273629 was >150-fold and
365-fold selective for iNOS against eNOS and nNOS, resp. The
pharmacokinetic profile of GW274150 was biphasic in healthy rats and mice
with a terminal half-life of .apprx.6 h. That of GW273629 was also
biphasic in rats, producing a terminal half-life of .apprx.3 h. In mice
however, elimination of GW273629 appeared monophasic and more rapid
(.apprx.10 min). Both compds. show a high oral bioavailability (>90%) in
rats and mice. GW274150 was effective in inhibiting LPS-induced plasma
NO_x levels in mice with an ED₅₀ of 3.2±0.7 mg kg⁻¹ after 14 h i.p. and
3.8±1.5 mg kg⁻¹ after 14 h when administered orally. GW274150 was
effective in inhibiting LPS-induced plasma NO_x levels in mice with an ED₅₀
of 3.2±0.7 mg kg⁻¹ after 14 h i.p. and 3.8±1.5 mg kg⁻¹ after 14 h
when administered orally. GW273629 showed shorter-lived effects on plasma
NO_x and an ED₅₀ of 9±2 mg kg⁻¹ after 2 h when administered i.p. The
effects of GW274150 and GW273629 in vivo were consistent with high
selectivity for iNOS, as these inhibitors were of low potency against nNOS
in the rat cerebellum and did not cause significant effects on blood
pressure in instrumented mice.

RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 2 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN
 AN 2005:48432 CAPLUS
 DN 142:169464
 TI Beneficial effects of GW274150 treatment on the development of
 experimental colitis induced by dinitrobenzene sulfonic acid
 AU Di Paola, Rosanna; Mazzon, Emanuela; Patel, Nimesh S. A.; Genovese,
 Tiziana; Muia, Carmelo; Thiemermann, Christoph; De Sarro, Angelina;
 Cuzzocrea, Salvatore
 CS Department of Clinical and Experimental Medicine and Pharmacology, School
 of Medicine, Policlinico Universitario, University of Messina Torre
 Biologica, Messina, 98123, Italy
 SO European Journal of Pharmacology (2005), 507(1-3), 281-289
 CODEN: EJPHAZ; ISSN: 0014-2999
 PB Elsevier B.V.
 DT Journal
 LA English
 AB Inflammatory bowel disease is associated with inducible nitric oxide synthase
 (iNOS) expression, oxidative and nitrosative stress, and leukocyte
 infiltration in the colon. Here, the authors investigate the effects of
 the selective iNOS-inhibitor (S)-2-amino-(1-iminoethylamino)-5-
 thiopentanoic acid (GW274150) on the development of exptl. colitis induced
 by dinitrobenzene sulfonic acid. When compared to dinitrobenzene sulfonic
 acid-treated mice, GW274150 (5 mg/kg i.p.)-treated mice subjected to
 dinitrobenzene sulfonic ACID-induced colitis experienced a significantly
 lower rate of the extent and severity of the histol. signs of colon
 injury. Dinitrobenzene sulfonic acid-treated mice experienced hemorrhagic
 diarrhea and weight loss. At 4 days after the administration of
 dinitrobenzene sulfonic acid, the mucosa of the colon exhibited large
 areas of necrosis. Immunohistochem. for nitrotyrosine and poly
 (ADP-ribose) (PAR) showed an intense staining in the inflamed colon.
 Treatment of dinitrobenzene sulfonic acid-treated mice with GW274150
 significantly reduced the degree of hemorrhagic diarrhea and weight loss
 caused by administration of dinitrobenzene sulfonic acid. GW274150 also
 caused a substantial reduction of (i) the degree of colon injury, (ii) the
 rise in myeloperoxidase (MPO) activity (mucosa), (iii) the increase in
 staining (immunohistochem.) for nitrotyrosine, as well as (iv) PARP
 activation caused by dinitrobenzene sulfonic acid in the colon. Thus,
 GW274150 treatment reduced the degree of colitis caused by dinitrobenzene
 sulfonic acid. The authors propose that selective inhibition of iNOS
 activity with GW274150 may be useful in the treatment of inflammatory
 bowel disease.
 RE.CNT 57 THERE ARE 57 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 3 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN
 AN 2004:346162 CAPLUS
 DN 140:399632
 TI Effects of GW274150, a novel and selective inhibitor of iNOS activity, in
 acute lung inflammation
 AU Dugo, Laura; Marzocco, Stefania; Mazzon, Emanuela; Di Paola, Rosanna;
 Genovese, Tiziana; Caputi, Achille P.; Cuzzocrea, Salvatore
 CS Department Clinical and Experimental Medicine and Pharmacology, University
 of Messina, Messina, 98100, Italy
 SO British Journal of Pharmacology (2004), 141(6), 979-987
 CODEN: BJPCBM; ISSN: 0007-1188
 PB Nature Publishing Group
 DT Journal
 LA English
 AB The aim of this study was to investigate the effect of GW274150, a novel,
 potent and selective inhibitor of inducible nitric oxide synthase (iNOS)
 activity in a model of lung injury induced by carrageenan administration
 in the rats. Injection of carrageenan into the pleural cavity of rats
 elicited an acute inflammatory response characterized by: fluid
 accumulation in the pleural cavity which contained a large number of

polymorphonuclear cells (PMNs) as well as an infiltration of PMNs in lung tissues and subsequent lipid peroxidn., and increased production of nitrite/nitrate (NOx), tumor necrosis factor α (TNF- α) and interleukin-1 β (IL-1 β). All parameters of inflammation were attenuated in a dose-dependent manner by GW274150 (2.5, 5 and 10 mg kg⁻¹ injected i.p. 5 min before carrageenan). Carrageenan induced an upregulation of the intracellular adhesion mols.-1 (ICAM-1), as well as nitrotyrosine and poly (ADP-ribose) (PAR) as determined by immunohistochem. anal. of lung tissues. The degree of staining for the ICAM-1, nitrotyrosine and PAR was reduced by GW274150. These results clearly confirm that NO from iNOS plays a role in the development of the inflammatory response by altering key components of the inflammatory cascade. GW274150 may offer a novel therapeutic approach for the management of various inflammatory diseases where NO and related radicals have been postulated to play a role.

RE.CNT 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 4 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN

AN 2004:2680 CAPLUS

DN 140:65201

TI (2S)-2-Amino-4-{[2-(ethanimidoylamino)ethyl]thio}butanoic acid nitric oxide synthase inhibitor in stabilized pharmaceutical dosage forms

IN Broughton, Stuart James; Gharu, Rajinder Kumar; Leow, Mark Yuon Tuck; Neale, Philip John

PA SB Pharmco Puerto Rico Inc., P. R.

SO PCT Int. Appl., 16 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004000296	A1	20031231	WO 2003-EP6465	20030619
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	EP 1513511	A1	20050316	EP 2003-740281	20030619
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
PRAI	GB 2002-14147	A	20020619		
	WO 2003-EP6465	W	20030619		
AB	Pharmaceutical compns. comprising (2S)-MeC(:NH)NHCH ₂ CH ₂ SCH ₂ CH ₂ CH(NH ₂)CO ₂ H (I) a pharmaceutically acceptable bulking agent and one or more antioxidants or chelating agents are described. A direct compression formula for tablets contained I, EDTA, Avical PH101, silica, and Mg stearate.				

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 5 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN

AN 2003:947089 CAPLUS

DN 140:314741

TI GW274150 inhibits nitric oxide production by primary cultures of rat proximal tubular cells

AU Chatterjee, Prabal K.; Kvale, Espen O.; Patel, Nimesh S. A.; Thiemermann, Christoph

CS Department of Experimental Medicine, Nephrology & Critical Care, William

Harvey Research Institute, Queen Mary - University of London, UK
SO Medical Science Monitor (2003), 9(10), BR357-BR362
CODEN: MSMOFR; ISSN: 1234-1010
PB International Scientific Literature, Inc.
DT Journal
LA English
AB Background: Production of nitric oxide (NO) subsequent to expression of inducible NO synthase (iNOS) contributes to the development of ischemic renal injury and inflammation. Here the authors investigate the effects of GW274150, a potent, long-acting and highly selective inhibitor of iNOS activity, on NO production by primary cultures of rat proximal tubular cells (PTC). Material/Methods: Pure populations of PTC were isolated from the cortex of kidneys obtained from male Wistar rats using a combination of collagenase digestion, sieving and Percoll centrifugation. Confluent PTC cultures were incubated for 1-24 h with MEM, interferon- γ (IFN- γ , 100 iu/mL), bacterial lipopolysaccharide (LPS, 10 μ g/mL) in combination after which NO production was determined PTC were also incubated with IFN- γ (100 iu/mL) and LPS (10 μ g/mL) and increasing concns. of GW274150 or L-N6-(1-iminoethyl)lysine (L-NIL) (10 nM - 1 mM) for 24 h after which nitrite levels in the incubation medium were measured. Results: IFN- γ and LPS in combination produced a significant, time-dependent increase in NO production Both GW274150 and L-NIL produced a significant and concentration-dependent inhibition of NO production However, GW274150 was markedly more potent (EC50 .apprx. 100 nM) than L-NIL (EC50 .apprx. 10 μ M). Conclusions: GW274150 inhibits NO production by primary cultures of PTCs and may therefore be useful in conditions associated with nitrosative stress of the kidney.
RE.CNT 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 6 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN
AN 2003:300915 CAPLUS
DN 138:302642
TI Inducible nitric oxide synthase inhibitors as vaccine adjuvants
IN Thomsen, Lindy Louise
PA Glaxo Group Limited, UK
SO PCT Int. Appl., 33 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003030935	A2	20030417	WO 2002-GB4365	20020926
	WO 2003030935	A3	20030814		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2462582	AA	20030417	CA 2002-2462582	20020926
	EP 1432440	A2	20040630	EP 2002-762572	20020926
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
	JP 2005510478	T2	20050421	JP 2003-533966	20020926
	US 2005054726	A1	20050310	US 2004-491843	20041011
PRAI	GB 2001-24022	A	20011005		
	WO 2002-GB4365	W	20020926		
OS	MARPAT 138:302642				

AB The present invention relates to the use of inducible nitric oxide synthase (iNOS) inhibitors as vaccine adjuvants, and in a preferred aspect of the invention they are used for adjuvanting nucleic acid (DNA) vaccines. The iNOS inhibitors preferably provide for an increase in antigen-specific CD4-pos. and/or CD8-pos. T cells. These compds. preferably induce a Th1-biased immune response as measured by increased formation of Th1 cytokines, in particular interferon γ . The present invention further provides pharmaceutical compns. comprising an antigen and the inhibitor.

L7 ANSWER 7 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN

AN 2003:289025 CAPLUS

DN 139:301665

TI GW274150, a potent and highly selective inhibitor of iNOS, reduces experimental renal ischemia/reperfusion injury

AU Chatterjee, Prabal K.; Patel, Nimesh S. A.; Sivarajah, Ahila; Kvale, Espen O.; Dugo, Laura; Cuzzocrea, Salvatore; Brown, Paul A. J.; Stewart, Keith N.; Mota-Filipe, Helder; Britti, Domenico; Yaqoob, Muhammad M.; Thiemermann, Christoph

CS Department of Experimental Medicine and Nephrology, The William Harvey Research Institute, Queen Mary, University of London, London, UK

SO Kidney International (2003), 63(3), 853-865

CODEN: KDYIA5; ISSN: 0085-2538

PB Blackwell Publishing, Inc.

DT Journal

LA English

AB Generation of nitric oxide (NO) by inducible nitric oxide synthase (iNOS) may contribute to renal ischemia/reperfusion (I/R) injury. The aim of this study was to investigate the effects of GW274150, a novel, highly selective, potent and long-acting inhibitor of iNOS activity in rat and mouse models of renal I/R. Rats were administered GW274150 (5 mg/kg i.v. bolus administered 30 min prior to I/R) and subjected to bilateral renal ischemia (45 min) followed by reperfusion (6 h). Serum and urinary indicators of renal dysfunction, tubular and reperfusion injury were measured, specifically, serum urea, creatinine, aspartate aminotransferase (AST) and N-acetyl- β -D-glucosaminidase (NAG) enzymuria. In addition, renal sections were used for histol. scoring of renal injury and for immunol. evidence of nitrotyrosine formation and poly [ADP (ADP)-ribose] (PAR). Nitrate levels were measured in rat plasma using the Griess assay. Mice (wild-type, administered 5 mg/kg GW274150, and iNOS-/-) were subjected to bilateral renal ischemia (30 min) followed by reperfusion (24 h) after which renal dysfunction (serum urea, creatinine), renal myeloperoxidase (MPO) activity and malondialdehyde (MDA) levels were measured. GW274150, administered prior to I/R, significantly reduced serum urea, serum creatinine, AST, and NAG indicating reduction of renal dysfunction and injury caused by I/R. GW274150 reduced histol. evidence of tubular injury and markedly reduced immunohistochem. evidence of nitrotyrosine and PAR formation, indicating reduced peroxynitrite formation and poly (ADP-ribose) polymerase (PARP) activation, resp. GW274150 abolished the rise in the plasma levels of nitrate (indicating reduced NO production). GW274150 also reduced the renal dysfunction in wild-type mice to levels similar to that observed in iNOS-/- mice subjected to I/R. Renal MPO activity and MDA levels were significantly reduced in wild-type mice administered GW274150 and iNOS-/- mice subjected to renal I/R, indicating reduced polymorphonuclear leukocyte (PMN) infiltration and lipid peroxidn. These results suggest that (1) an enhanced formation of NO by iNOS contributes to the pathophysiol. of renal I/R injury and (2) GW274150 reduces I/R injury of the kidney. We propose that selective inhibitors of iNOS activity may be useful against renal dysfunction and injury associated with I/R of the kidney.

RE.CNT 67 THERE ARE 67 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 8 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN

AN 2003:51504 CAPLUS

DN 139:159864
TI A novel, potent and selective inhibitor of the activity of inducible nitric oxide synthase (GW274150) reduces the organ injury in hemorrhagic shock
AU McDonald, M. C.; Izumi, M.; Cuzzocrea, S.; Thiemermann, C.
CS The William Harvey Research Institute, St. Bartholomew's and The Royal London School of Medicine and Dentistry, London, EC1M6BQ, UK
SO Journal of Physiology and Pharmacology (2002), 53(4, Pt. 1), 555-569
CODEN: JPHPEI; ISSN: 0867-5910
PB Polish Physiological Society
DT Journal
LA English
AB An enhanced formation of nitric oxide (NO) by the inducible NO synthase (iNOS) may contribute to the pathophysiol. of hemorrhagic shock. This study investigates the effect of a novel, potent and selective inhibitor of iNOS activity (GW274150) on the circulatory failure and the organ injury and dysfunction associated with hemorrhagic shock in the anesthetized rat. Hemorrhage (sufficient to lower mean arterial blood pressure to 45 mmHg for 90 min) and subsequent resuscitation with shed blood resulted (within 4 h after resuscitation) in a delayed fall in blood pressure, renal and liver injury and dysfunction as well as the pancreatic injury. Pre-treatment of rats with GW274150 (5 mg/kg at 30 min prior to the onset of hemorrhage) attenuated the renal dysfunction as well as the liver and pancreatic injury caused by hemorrhage and resuscitation. Interestingly, GW274150 did not reduce the delayed fall in blood pressure associated with hemorrhagic shock. We propose that an enhanced formation of NO from iNOS contributes to the organ injury and dysfunction in hemorrhagic shock, and that highly selective inhibitors of iNOS activity, such as GW274150, may represent a novel therapeutic approach for the therapy of hemorrhagic shock.

RE.CNT 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 9 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN
AN 2002:804285 CAPLUS
DN 138:314136
TI Beneficial effects of GW274150, a novel, potent and selective inhibitor of iNOS activity, in a rodent model of collagen-induced arthritis
AU Cuzzocrea, Salvatore; Chatterjee, Prabal K.; Mazzon, Emanuela; McDonald, Michelle C.; Dugo, Laura; Di Paola, Rosanna; Serraino, Ivana; Britti, Domenico; Caputi, Achille P.; Thiemermann, Christoph
CS School of Medicine, Institute of Pharmacology, University of Messina, Policlinico Universitario, Gazzi, Messina, 98100, Italy
SO European Journal of Pharmacology (2002), 453(1), 119-129
CODEN: EJPHAZ; ISSN: 0014-2999
PB Elsevier Science B.V.
DT Journal
LA English
AB The aim of this study was to investigate the role of inducible nitric oxide synthase (iNOS) on the modulation of the inflammatory response in mice subjected to collagen-induced arthritis. Collagen-induced arthritis was induced in wild-type mice (iNOS-WT) treated with GW274150, a novel, potent and selective inhibitor of iNOS activity, and in mice lacking the gene for iNOS (iNOS knock-out', iNOS-KO), by an intradermal injection of 100 µl of emulsion containing 100 µg of bovine type II collagen and complete Freund's adjuvant at the base of the tail. After 21 days, a second injection of type II collagen in complete Freund's adjuvant was administered. iNOS-WT mice developed erosive hind paw arthritis when immunized with type II collagen in complete Freund's adjuvant. Over a 35-day period, macroscopic clin. evidence of collagen-induced arthritis first appeared as periarticular erythema and edema in the hind paws. By day 28, the incidence of collagen-induced arthritis was 100% in type II collagen-challenged iNOS-WT mice and the severity of collagen-induced arthritis progressed with radiog. evaluation revealing resorption of bone. Histopathol. of collagen-induced arthritis mice demonstrated erosion of

the cartilage at the joint margins. iNOS-WT mice treated with GW274150 (5 mg/kg, i.p. daily) starting at the onset of arthritis (day 23), and iNOS-KO mice showed a delay of the development of the clin. signs at days 24-35 and an improvement of the histol. status in the knee and paw. Immunohistochem. anal. for nitrotyrosine and for poly(ADP-ribose) polymerase revealed pos. staining in inflamed joints from type II collagen-treated iNOS-WT mice. The degree of staining for nitrotyrosine and poly(ADP-ribose) polymerase were markedly reduced in tissue sections obtained from type II collagen-treated iNOS-WT mice, who had received GW274150 and from iNOS-KO mice. Furthermore, radiog. signs of protection against bone resorption were present in the joints of iNOS-WT mice treated with GW274150 as well as in the joint from iNOS-KO mice. This study provides the first evidence that GW274150, a novel, potent and selective inhibitor of iNOS activity, attenuates the degree of chronic inflammation and tissue damage associated with collagen-induced arthritis in mice. Furthermore, these results suggest that the induction of iNOS and NO production are essential for the up-regulation of the inflammatory response during exptl. collagen-induced arthritis.

RE.CNT 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN
AN 2001:597331 CAPLUS
DN 136:288829
TI GW274150 is a potent, long-acting, highly-selective inhibitor of inducible nitric oxide synthase (NOS-2) with therapeutic potential in post-operative ileus
AU Alderton, W.; Angell, A.; Clayton, N.; Craig, C.; Dawson, J.; Frend, A.; McGill, J.; Mangel, A.; Moncada, S.; Rees, D.; Russell, L.; Schwartz, S.; Waslidge, N.; Knowles, R.
CS Glaxo Wellcome R and D, Stevenage, SG1 2NY, UK
SO Portland Press Proceedings (2000), 16(Biology of Nitric Oxide, Part 7), 22
CODEN: POPPEF; ISSN: 0966-4068
PB Portland Press Ltd.
DT Journal
LA English
AB GW274150 [(S)-2-amino-7-acetamidino-5-thioheptanoic acid] is a novel α -amino acid that potently inhibited human inducible nitric oxide synthase (iNOS) with selectivity vs. human eNOS and nNOS. In studies with purified NOS isoforms, GW274150 was a time-dependent, arginine-site inhibitor of iNOS and a rapidly-reversible inhibitor of eNOS. This novel compound had a long pharmacokinetic half-life and high oral bioavailability in several species. The selectivity of GW274150 against the constitutive NOS isoforms was maintained in vivo, the compound producing no significant effect on conscious mouse blood pressure dosed at 100 mg/kg and on rat brain plus nitrite levels at 50 mg/kg. Post-operative ileus is one potential therapeutic application for GW274150. In a rat model of post-operative ileus, GW274150 was maximally effective at 1-5 mg/kg, yielding a 67% reversal of delayed GI transit. The compound was also effective in a rat model of acute inflammatory pain (adjuvant).

RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 11 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN
AN 2000:209102 CAPLUS
DN 133:12344
TI Inhibition of inducible nitric oxide synthase by acetamidine derivatives of hetero-substituted lysine and homolysine
AU Young, Robert J.; Beams, Richard M.; Carter, Keith; Clark, Helen A. R.; Coe, Diane M.; Chambers, C. Lynn; Davies, P. Ifeyinwa; Dawson, John; Drysdale, Martin J.; Franzman, Karl W.; French, Colin; Hodgson, Simon T.; Hodson, Harold F.; Kleanthous, Savvas; Rider, Peter; Sanders, Daniela; Sawyer, David A.; Scott, Keith J.; Shearer, Barry G.; Stocker, Richard; Smith, Steven; Tackley, Miriam C.; Knowles, Richard G.
CS Glaxo Wellcome Research and Development, Stevenage, SG1 2NY, UK

SO Bioorganic & Medicinal Chemistry Letters (2000), 10(6), 597-600
 CODEN: BMCLE8; ISSN: 0960-894X
 PB Elsevier Science Ltd.
 DT Journal
 LA English
 AB The synthesis and in vitro evaluation of the acetamidine derivs. of hetero-substituted lysine and homolysine analogs have identified potent inhibitors of human nitric oxide synthase enzymes, including examples with marked selectivity for the inducible isoform.
 RE.CNT 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 12 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN
 AN 1999:753054 CAPLUS
 DN 131:346497
 TI Use of nitric oxide synthase inhibitors in the manufacture of a medicament for the prophylaxis or treatment of bacterial infection
 IN Alderton, Wendy Karen; Knowles, Richard Graham; Ladel, Christoph Hubertus
 PA Glaxo Group Limited, UK
 SO PCT Int. Appl., 34 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9959566	A1	19991125	WO 1999-EP3265	19990512
	W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	AU 9940406	A1	19991206	AU 1999-40406	19990512
PRAI	GB 1998-10299	A	19980515		
	WO 1999-EP3265	W	19990512		

OS MARPAT 131:346497
 AB Inducible nitric oxide synthase inhibitors are used for the manufacture of a medicament for the prophylaxis or treatment of a bacterial infection, where the inhibitor of inducible nitric oxide synthase is e.g.
 HN:C(R1)NHR2 [R1 = C1-6 straight or branched chain alkyl; Q = QC(NH2)CO2H (Q = alkylene, alkenylene, etc.), ring-substituted benzyl] or a pharmaceutically acceptable salt, ester, or amide thereof.
 RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2005 ACS on STN
 AN 1998:490618 CAPLUS
 DN 129:122862
 TI Preparation of S-[2-(1-iminoethylamino)ethyl]homocysteine as nitric oxide synthase inhibitor
 IN Beams, Richard Mansfield; Drysdale, Martin James; Franzman, Karl Witold; Friend, Anthony Joseph; Hodson, Harold Francis; Knowles, Richard Graham; Rees, Daryl David; Sawyer, David Alan
 PA Glaxo Group Ltd., UK; Beams, Richard Mansfield; Drysdale, Martin James; Franzman, Karl Witold; Friend, Anthony Joseph; Hodson, Harold Francis; Knowles, Richard Graham; Rees, Daryl David; Sawyer, David Alan
 SO PCT Int. Appl., 29 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9830537	A1	19980716	WO 1998-EP96	19980109
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	CA 2277877	AA	19980716	CA 1998-2277877	19980109
	AU 9862083	A1	19980803	AU 1998-62083	19980109
	AU 723095	B2	20000817		
	ZA 9800179	A	19990709	ZA 1998-179	19980109
	EP 958277	A1	19991124	EP 1998-904050	19980109
	EP 958277	B1	20011121		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	EE 9900281	A	20000215	EE 1999-281	19980109
	EE 4013	B1	20030415		
	JP 2000504041	T2	20000404	JP 1998-530549	19980109
	JP 3251301	B2	20020128		
	BR 9806870	A	20000418	BR 1998-6870	19980109
	NZ 336379	A	20010126	NZ 1998-336379	19980109
	AT 209183	E	20011215	AT 1998-904050	19980109
	PT 958277	T	20020531	PT 1998-904050	19980109
	ES 2168737	T3	20020616	ES 1998-904050	19980109
	SK 283201	B6	20030304	SK 1999-933	19980109
	AP 1204	A	20030915	AP 1999-1603	19980109
	W: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW				
	IL 130551	A1	20040104	IL 1998-130551	19980109
	CZ 293099	B6	20040218	CZ 1999-2483	19980109
	TW 502010	B	20020911	TW 1998-87100434	19980114
	TW 538021	B	20030621	TW 1999-88103866	19980114
	NO 9903429	A	19990712	NO 1999-3429	19990712
	NO 312192	B1	20020408		
	US 6369272	B1	20020409	US 1999-341220	19990824
	HK 1021531	A1	20020315	HK 2000-100440	20000124
	US 2002010366	A1	20020124	US 2001-930605	20010815
	US 6620848	B2	20030916		
PRAI	US 1997-69882P	P	19970113		
	US 1997-783402	A	19970113		
	WO 1998-EP96	W	19980109		
	US 1999-341220	A1	19990824		
OS	MARPAT 129:122862				
AB	HN:CM ₂ NHCH ₂ CH ₂ SCH ₂ CH ₂ CH(NH ₂)CO ₂ H (I) was prepared for use as a selective inhibitor of nitric oxide synthase (NOS). Thus, (S)-I was prepared by treatment of L-homocystine with Na in liquid NH ₃ and then N-benzyloxycarbonylethanolamine tosylate, cleavage of the benzyloxycarbonyl protecting group with HBr in AcOH, and reaction with Et acetimidate hydrochloride. (S)-I was assayed for inhibition of inducible and endothelial NOS (IC ₅₀ = 0.73 and 43 μM, resp.).				
RE.CNT	7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT				

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	59.41	114.30
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-10.95	-10.95

STN INTERNATIONAL LOGOFF AT 15:01:11 ON 22 AUG 2005

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:ssptamxgl614

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 FEB 27 New STN AnaVist pricing effective March 1, 2006
NEWS 4 MAY 10 CA/CAPLUS enhanced with 1900-1906 U.S. patent records
NEWS 5 MAY 11 KOREAPAT updates resume
NEWS 6 MAY 19 Derwent World Patents Index to be reloaded and enhanced
NEWS 7 MAY 30 IPC 8 Rolled-up Core codes added to CA/CAPLUS and
USPATFULL/USPAT2
NEWS 8 MAY 30 The F-Term thesaurus is now available in CA/CAPLUS
NEWS 9 JUN 02 The first reclassification of IPC codes now complete in
INPADOC
NEWS 10 JUN 26 TULSA/TULSA2 reloaded and enhanced with new search and
and display fields
NEWS 11 JUN 28 Price changes in full-text patent databases EPFULL and PCTFULL
NEWS 12 JUL 11 CHEMSAFE reloaded and enhanced
NEWS 13 JUL 14 FSTA enhanced with Japanese patents
NEWS 14 JUL 19 Coverage of Research Disclosure reinstated in DWPI
NEWS 15 AUG 09 INSPEC enhanced with 1898-1968 archive
NEWS 16 AUG 28 ADISCTI Reloaded and Enhanced
NEWS 17 AUG 30 CA(SM)/CAPLUS(SM) Austrian patent law changes
NEWS 18 SEP 11 CA/CAPLUS enhanced with more pre-1907 records

NEWS EXPRESS JUNE 30 CURRENT WINDOWS VERSION IS V8.01b, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 26 JUNE 2006.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8
NEWS X25 X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 07:59:17 ON 19 SEP 2006

=> file caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'CAPLUS' ENTERED AT 07:59:27 ON 19 SEP 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 Sep 2006 VOL 145 ISS 13

FILE LAST UPDATED: 18 Sep 2006 (20060918/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s 210354-22-6/rn

16 210354-22-6

1 210354-22-6D

L1 16 210354-22-6/RN

(210354-22-6 (NOTL) 210354-22-6D)

=> d 1-16 bib abs hitstr

L1 ANSWER 1 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2006:315097 CAPLUS

DN 145:296

TI Design, synthesis, and evaluation of new type of L-amino acids containing pyridine moiety as nitric oxide synthase inhibitor

AU Ijuin, Ryosuke; Umezawa, Naoki; Higuchi, Tsunehiko

CS Graduate School of Pharmaceutical Sciences, Nagoya City University, Nagoya, 467-8603, Japan

SO Bioorganic & Medicinal Chemistry (2006), 14(10), 3563-3570


CODEN: BMECEP; ISSN: 0968-0896

PB Elsevier B.V.

DT Journal

LA English

GI



I

2 HCl


AB New amino acids were designed and synthesized as candidate inhibitors of human nitric oxide synthase (NOS). The 2-aminopyridine-containing L-amino acids I had potent inhibitory activity toward all of the human NOS isoenzymes. A computational docking study was carried out to investigate the mechanism of the inhibitory effect.

IT 210354-22-6, GW 274150
 RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (pyridyl amino acids as nitric oxide synthase inhibitors)

RN 210354-22-6 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 2 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2006:56531 CAPLUS

DN 145:40027

TI Role of inducible nitric oxide synthase in the reduced responsiveness of the myocardium to catecholamines in a hyperdynamic, murine model of septic shock

AU Barth, Eberhard; Radermacher, Peter; Thiernemann, Christoph; Weber, Sandra; Georgieff, Michael; Albuszies, Gerd

CS Sektion Anaesthesiologische Pathophysiologie und Verfahrensentwicklung, Universitaetsklinikum, Ulm, Germany

SO Critical Care Medicine (2006), 34(2), 307-313
 CODEN: CCMDC7; ISSN: 0090-3493

PB Lippincott Williams & Wilkins

DT Journal

LA English

AB Objectives: Excess nitric oxide production is a key mediator of hypotension and catecholamine-resistance in septic shock. Although nitric oxide synthase blockade has been shown to restore hemodynamics, conflicting results on myocardial function were reported. Inducible nitric oxide synthase (iNOS) knockout (iNOS-/-) mice showed improved heart function, but these results were obtained during hypodynamic shock characterized by reduced cardiac output. Therefore, we investigated heart function and

catecholamine responsiveness in a clin. relevant, murine model of cecal ligation and puncture (CLP)-induced septic shock. Design: Prospective, controlled, randomized animal study. Setting: University animal research laboratory Subjects: Male C57Bl/6 wild-type and iNOS-/- mice. Interventions: Fifteen hours after CLP, three groups of mice (wild-type controls, n = 9; iNOS-/-, n = 12; and wild-type mice receiving 5 mg·kg⁻¹ i.p. of the selective iNOS inhibitor GW274150 immediately after CLP, n = 8) were anesthetized, mech. ventilated, and instrumented (central venous and left ventricular pressure-conductance catheter). Measurements were recorded 18, 21, and 24 h post-CLP. Hydroxyethylstarch and norepinephrine were infused to achieve normotensive and hyperdynamic hemodynamics. Measurements and main results: There was no intergroup difference in mean arterial pressure, stroke volume, and left ventricular ejection fraction. Norepinephrine doses required to achieve the hemodynamic targets were lower in GW274150 (p < .001 vs. controls) and even further reduced in iNOS-/- mice (p < .001 vs. controls, p < .001 vs. GW274150). In the control group, the higher norepinephrine doses resulted in significantly higher heart rates and consequently cardiac output, maximal contraction, and relaxation than in the GW274150 and iNOS-/- animals. Left ventricular end-diastolic volume was also significantly higher in the controls than in the GW274150 and iNOS-/- mice, whereas left ventricular end-diastolic pressure did not differ. Conclusions: Our results confirm septic shock-related impaired left ventricular function. Genetic iNOS deletion and pharmacol. iNOS blockade enhanced cardiac norepinephrine responsiveness due to improved systolic function. In contrast, iNOS inhibition seemed to be affiliated with compromised left ventricular relaxation.

IT 210354-22-6, GW274150


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(inducible nitric oxide synthase inhibitor GW274150 enhanced myocardium responsiveness to catecholamine by improving systolic function and thus maintained left ventricular function in hyperdynamic murine model of septic shock)

RN 210354-22-6 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 3 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2006:35943 CAPLUS

DN 145:20904

TI GW274150, a novel and highly selective inhibitor of the inducible isoform of nitric oxide synthase (iNOS), shows analgesic effects in rat models of inflammatory and neuropathic pain

AU De Alba, Jorge; Clayton, Nick M.; Collins, Sue D.; Colthup, Philip; Chessell, Iain; Knowles, Richard G.

CS Department of Respiratory Pharmacology, RI CEDD GlaxoSmithKline Research and Development, Medicines Research Centre, Hertfordshire, SG1 2NY, UK

SO Pain (2006), 120(1-2), 170-181

CODEN: PAINDB; ISSN: 0304-3959

PB Elsevier Ltd.

DT Journal

LA English

AB Nitric oxide (NO), synthesized by different isoforms of nitric oxide synthase (NOS), has been linked with the development and maintenance of nociception. We studied the role of the inducible isoform, iNOS, in two different rat pain models with an inflammatory component. iNOS was immunohistochem. detected locally in the paw 6 h after Freund's Complete Adjuvant (FCA) injection, showing a plateau at 24-72 h and falling slowly in the following weeks. This correlated with the late phase of the hypersensitivity to pain revealed in the behavioral tests. A highly selective iNOS inhibitor GW274150 (1-30 mg/kg orally, 24 h after FCA) suppressed the accumulation of nitrite in the inflamed paw indicating substantial iNOS inhibition. At the same time it partially reversed FCA-induced hypersensitivity to pain and edema in a dose-dependent manner. After Chronic Constriction Injury (CCI) surgery to the sciatic nerve, iNOS presence was only detected locally in the region of the nerve (inflammatory cells). GW274150 (3-30 mg/kg orally, 21 days after surgery) also reversed significantly the CCI-associated hypersensitivity to pain. No iNOS was detectable in dorsal root ganglia, spinal cord or brain in either model. This study demonstrates a role for peripherally-expressed iNOS in pain conditions with an inflammatory component and the potential value of iNOS inhibitors in such conditions.

IT 210354-22-6, GW274150


RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); PKT (Pharmacokinetics); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(iNOS inhibitor GW274150 suppressed nitrite accumulation in inflamed paw, partially reversed FCA-induced hypersensitivity to pain and edema in dose-dependent manner, significantly reversed CCI-associated hypersensitivity to pain in rat model)

RN 210354-22-6 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 73 THERE ARE 73 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 4 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2005:465475 CAPLUS

DN 143:71325

TI GW274150 and GW273629 are potent and highly selective inhibitors of inducible nitric oxide synthase in vitro and in vivo

AU Alderton, Wendy K.; Angell, Anthony D. R.; Craig, Caroline; Dawson, John; Garvey, Edward; Moncada, Salvador; Monkhouse, Jayne; Rees, Daryl; Russell, Linda J.; Russell, Rachel J.; Schwartz, Sheila; Waslidge, Neil; Knowles, Richard G.

CS Medicines Research Centre, Respiratory & Inflammation Centre of Excellence for Drug Discovery, GlaxoSmithKline Research, Stevenage, SG1 2NY, UK

SO British Journal of Pharmacology (2005), 145(3), 301-312

CODEN: BJPCBM; ISSN: 0007-1188

PB Nature Publishing Group

DT Journal


LA English

AB GW274150 ([2-[(1-iminoethyl) amino]ethyl]-L-homocysteine) and GW273629 (3-[[2-[(1-iminoethyl)amino]ethyl]sulfonyl]-L-alanine) are potent, time-dependent, highly selective inhibitors of human inducible nitric oxide synthase (iNOS) vs. endothelial NOS (eNOS) (>100-fold) or neuronal NOS (nNOS) (>80-fold). GW274150 and GW273629 are arginine competitive,

NADPH-dependent inhibitors of human iNOS with steady state K_d values of <40 and <90 nM, resp. GW274150 and GW273629 inhibited intracellular iNOS in J774 cells in a time-dependent manner, reaching IC₅₀ values of 0.2±0.04 and 1.3±0.16 µM, resp. They were also acutely selective in intact rat tissues: GW274150 was >260-fold and 219-fold selective for iNOS against eNOS and nNOS, resp., while GW273629 was >150-fold and 365-fold selective for iNOS against eNOS and nNOS, resp. The pharmacokinetic profile of GW274150 was biphasic in healthy rats and mice with a terminal half-life of .apprx.6 h. That of GW273629 was also biphasic in rats, producing a terminal half-life of .apprx.3 h. In mice however, elimination of GW273629 appeared monophasic and more rapid (.apprx.10 min). Both compds. show a high oral bioavailability (>90%) in rats and mice. GW274150 was effective in inhibiting LPS-induced plasma NO_x levels in mice with an ED₅₀ of 3.2±0.7 mg kg⁻¹ after 14 h i.p. and 3.8±1.5 mg kg⁻¹ after 14 h when administered orally. GW274150 was effective in inhibiting LPS-induced plasma NO_x levels in mice with an ED₅₀ of 3.2±0.7 mg kg⁻¹ after 14 h i.p. and 3.8±1.5 mg kg⁻¹ after 14 h when administered orally. GW273629 showed shorter-lived effects on plasma NO_x and an ED₅₀ of 9±2 mg kg⁻¹ after 2 h when administered i.p. The effects of GW274150 and GW273629 in vivo were consistent with high selectivity for iNOS, as these inhibitors were of low potency against nNOS in the rat cerebellum and did not cause significant effects on blood pressure in instrumented mice.

IT 210354-22-6, GW274150
 RL: PAC (Pharmacological activity); PKT (Pharmacokinetics); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (GW274150 and GW273629 are potent and highly selective inhibitors of inducible nitric oxide synthase in vitro and in vivo)
 RN 210354-22-6 CAPLUS
 CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 5 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2005:48432 CAPLUS
 DN 142:169464
 TI Beneficial effects of GW274150 treatment on the development of experimental colitis induced by dinitrobenzene sulfonic acid
 AU Di Paola, Rosanna; Mazzon, Emanuela; Patel, Nimesh S. A.; Genovese, Tiziana; Muia, Carmelo; Thiemermann, Christoph; De Sarro, Angelina; Cuzzocrea, Salvatore
 CS Department of Clinical and Experimental Medicine and Pharmacology, School of Medicine, Policlinico Universitario, University of Messina Torre Biologica, Messina, 98123, Italy
 SO European Journal of Pharmacology (2005), 507(1-3), 281-289
 CODEN: EJPHAZ; ISSN: 0014-2999
 PB Elsevier B.V.
 DT Journal
 LA English
 AB Inflammatory bowel disease is associated with inducible nitric oxide synthase (iNOS) expression, oxidative and nitrosative stress, and leukocyte infiltration in the colon. Here, the authors investigate the effects of the selective iNOS-inhibitor (S)-2-amino-(1-iminoethylamino)-5-thiopentanoic acid (GW274150) on the development of exptl. colitis induced

by dinitrobenzene sulfonic acid. When compared to dinitrobenzene sulfonic acid-treated mice, GW274150 (5 mg/kg i.p.)-treated mice subjected to dinitrobenzene sulfonic ACID-induced colitis experienced a significantly lower rate of the extent and severity of the histol. signs of colon injury. Dinitrobenzene sulfonic acid-treated mice experienced hemorrhagic diarrhea and weight loss. At 4 days after the administration of dinitrobenzene sulfonic acid, the mucosa of the colon exhibited large areas of necrosis. Immunohistochem. for nitrotyrosine and poly (ADP-ribose) (PAR) showed an intense staining in the inflamed colon. Treatment of dinitrobenzene sulfonic acid-treated mice with GW274150 significantly reduced the degree of hemorrhagic diarrhea and weight loss caused by administration of dinitrobenzene sulfonic acid. GW274150 also caused a substantial reduction of (i) the degree of colon injury, (ii) the rise in myeloperoxidase (MPO) activity (mucosa), (iii) the increase in staining (immunohistochem.) for nitrotyrosine, as well as (iv) PARP activation caused by dinitrobenzene sulfonic acid in the colon. Thus, GW274150 treatment reduced the degree of colitis caused by dinitrobenzene sulfonic acid. The authors propose that selective inhibition of iNOS activity with GW274150 may be useful in the treatment of inflammatory bowel disease.

IT 210354-22-6, GW274150
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (beneficial effects of GW274150 treatment on development of exptl. colitis induced by dinitrobenzene sulfonic acid)
 RN 210354-22-6 CAPLUS
 CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 57 THERE ARE 57 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 6 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2004:346162 CAPLUS
 DN 140:399632
 TI Effects of GW274150, a novel and selective inhibitor of iNOS activity, in acute lung inflammation
 AU Dugo, Laura; Marzocco, Stefania; Mazzon, Emanuela; Di Paola, Rosanna; Genovese, Tiziana; Caputi, Achille P.; Cuzzocrea, Salvatore
 CS Department Clinical and Experimental Medicine and Pharmacology, University of Messina, Messina, 98100, Italy
 SO British Journal of Pharmacology (2004), 141(6), 979-987
 CODEN: BJPCBM; ISSN: 0007-1188
 PB Nature Publishing Group
 DT Journal
 LA English
 AB The aim of this study was to investigate the effect of GW274150, a novel, potent and selective inhibitor of inducible nitric oxide synthase (iNOS) activity in a model of lung injury induced by carrageenan administration in the rats. Injection of carrageenan into the pleural cavity of rats elicited an acute inflammatory response characterized by: fluid accumulation in the pleural cavity which contained a large number of polymorphonuclear cells (PMNs) as well as an infiltration of PMNs in lung tissues and subsequent lipid peroxidn., and increased production of nitrite/nitrate (NOx), tumor necrosis factor α (TNF- α) and interleukin-1 β (IL-1 β). All parameters of inflammation were

attenuated in a dose-dependent manner by GW274150 (2.5, 5 and 10 mg kg⁻¹ injected i.p. 5 min before carrageenan). Carrageenan induced an upregulation of the intracellular adhesion mols.-1 (ICAM-1), as well as nitrotyrosine and poly (ADP-ribose) (PAR) as determined by immunohistochem. anal. of lung tissues. The degree of staining for the ICAM-1, nitrotyrosine and PAR was reduced by GW274150. These results clearly confirm that NO from iNOS plays a role in the development of the inflammatory response by altering key components of the inflammatory cascade. GW274150 may offer a novel therapeutic approach for the management of various inflammatory diseases where NO and related radicals have been postulated to play a role.

IT 210354-22-6, GW274150


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(effects of GW274150, novel and selective inhibitor of iNOS activity, in acute lung inflammation)

RN 210354-22-6 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 7 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2004:2680 CAPLUS

DN 140:65201

TI (2S)-2-Amino-4-{[2-(ethanimidoethylamino)ethyl]thio}butanoic acid nitric oxide synthase inhibitor in stabilized pharmaceutical dosage forms

IN Broughton, Stuart James; Gharu, Rajinder Kumar; Leow, Mark Yuen Tuck; Neale, Philip John

PA SB Pharmco Puerto Rico Inc., P. R.

SO PCT Int. Appl., 16 pp.

CODEN: PIXXD2

DT Patent


LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2004000296	A1	20031231	WO 2003-EP6465	20030619
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	AU 2003278958	A1	20040106	AU 2003-278958	20030619
	EP 1513511	A1	20050316	EP 2003-740281	20030619
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
	JP 2005533075	T2	20051104	JP 2004-514780	20030619
	US 2005222260	A1	20051006	US 2004-517801	20041210
PRAI	GB 2002-14147	A	20020619		

WO 2003-EP6465 W 20030619
 AB Pharmaceutical compns. comprising (2S)-MeC(=NH)NHCH₂CH₂SCH₂CH₂CH(NH₂)CO₂H (I) a pharmaceutically acceptable bulking agent and one or more antioxidants or chelating agents are described. A direct compression formula for tablets contained I, EDTA, Avical PH101, silica, and Mg stearate.
 IT 210354-22-6
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 ((2S)-2-Amino-4-{[2-(ethanimidoylamino)ethyl]thio}butanoic acid nitric oxide synthase inhibitor in stabilized pharmaceutical dosage forms)
 RN 210354-22-6 CAPLUS
 CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT


L1 ANSWER 8 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 2003:947089 CAPLUS
 DN 140:314741
 TI GW274150 inhibits nitric oxide production by primary cultures of rat proximal tubular cells
 AU Chatterjee, Prabal K.; Kvale, Espen O.; Patel, Nimesh S. A.; Thiemermann, Christoph
 CS Department of Experimental Medicine, Nephrology & Critical Care, William Harvey Research Institute, Queen Mary - University of London, UK
 SO Medical Science Monitor (2003), 9(10), BR357-BR362
 CODEN: MSMOFR; ISSN: 1234-1010
 PB International Scientific Literature, Inc.
 DT Journal
 LA English
 AB Background: Production of nitric oxide (NO) subsequent to expression of inducible NO synthase (iNOS) contributes to the development of ischemic renal injury and inflammation. Here the authors investigate the effects of GW274150, a potent, long-acting and highly selective inhibitor of iNOS activity, on NO production by primary cultures of rat proximal tubular cells (PTC). Material/Methods: Pure populations of PTC were isolated from the cortex of kidneys obtained from male Wistar rats using a combination of collagenase digestion, sieving and Percoll centrifugation. Confluent PTC cultures were incubated for 1-24 h with MEM, interferon- γ (IFN- γ , 100 iu/mL), bacterial lipopolysaccharide (LPS, 10 μ g/mL) in combination after which NO production was determined PTC were also incubated with IFN- γ (100 iu/mL) and LPS (10 μ g/mL) and increasing concns. of GW274150 or L-N6-(1-iminoethyl)lysine (L-NIL) (10 nM - 1 mM) for 24 h after which nitrite levels in the incubation medium were measured. Results: IFN- γ and LPS in combination produced a significant, time-dependent increase in NO production Both GW274150 and L-NIL produced a significant and concentration-dependent inhibition of NO production However, GW274150 was markedly more potent (EC50 .apprx. 100 nM) than L-NIL (EC50 .apprx. 10 μ M). Conclusions: GW274150 inhibits NO production by primary cultures of PTCs and may therefore be useful in conditions associated with nitrosative stress of the kidney.
 IT 210354-22-6, GW274150
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(GW274150 inhibits nitric oxide production by primary cultures of rat proximal tubular cells incubated with interferon- γ and LPS)

RN 210354-22-6 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 9 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2003:300915 CAPLUS

DN 138:302642

TI Inducible nitric oxide synthase inhibitors as vaccine adjuvants

IN Thomsen, Lindy Louise

PA Glaxo Group Limited, UK

SO PCT Int. Appl., 33 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003030935	A2	20030417	WO 2002-GB4365	20020926
	WO 2003030935	A3	20030814		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	CA 2462582	AA	20030417	CA 2002-2462582	20020926
	EP 1432440	A2	20040630	EP 2002-762572	20020926
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
	JP 2005510478	T2	20050421	JP 2003-533966	20020926
	US 2005054726	A1	20050310	US 2004-491843	20041011
PRAI	GB 2001-24022	A	20011005		
	WO 2002-GB4365	W	20020926		

OS MARPAT 138:302642

AB The present invention relates to the use of inducible nitric oxide synthase (iNOS) inhibitors as vaccine adjuvants, and in a preferred aspect of the invention they are used for adjuvanting nucleic acid (DNA) vaccines. The iNOS inhibitors preferably provide for an increase in antigen-specific CD4-pos. and/or CD8-pos. T cells. These compds. preferably induce a Th1-biased immune response as measured by increased formation of Th1 cytokines, in particular interferon γ . The present invention further provides pharmaceutical compns. comprising an antigen and the inhibitor.


IT 210354-22-6, GW 274150

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(inducible nitric oxide synthase inhibitors as vaccine adjuvants)

RN 210354-22-6 CAPLUS
CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L1 ANSWER 10 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2003:289025 CAPLUS
DN 139:301665
TI GW274150, a potent and highly selective inhibitor of iNOS, reduces experimental renal ischemia/reperfusion injury
AU Chatterjee, Prabal K.; Patel, Nimesh S. A.; Sivarajah, Ahila; Kvale, Espen O.; Dugo, Laura; Cuzzocrea, Salvatore; Brown, Paul A. J.; Stewart, Keith N.; Mota-Filipe, Helder; Britti, Domenico; Yaqoob, Muhammad M.; Thiernemann, Christoph
CS Department of Experimental Medicine and Nephrology, The William Harvey Research Institute, Queen Mary, University of London, London, UK
SO Kidney International (2003), 63(3), 853-865
CODEN: KDYIA5; ISSN: 0085-2538
PB Blackwell Publishing, Inc.
DT Journal
LA English
AB Generation of nitric oxide (NO) by inducible nitric oxide synthase (iNOS) may contribute to renal ischemia/reperfusion (I/R) injury. The aim of this study was to investigate the effects of GW274150, a novel, highly selective, potent and long-acting inhibitor of iNOS activity in rat and mouse models of renal I/R. Rats were administered GW274150 (5 mg/kg i.v. bolus administered 30 min prior to I/R) and subjected to bilateral renal ischemia (45 min) followed by reperfusion (6 h). Serum and urinary indicators of renal dysfunction, tubular and reperfusion injury were measured, specifically, serum urea, creatinine, aspartate aminotransferase (AST) and N-acetyl-β-D-glucosaminidase (NAG) enzymuria. In addition, renal sections were used for histol. scoring of renal injury and for immunol. evidence of nitrotyrosine formation and poly [ADP (ADP)-ribose] (PAR). Nitrate levels were measured in rat plasma using the Griess assay. Mice (wild-type, administered 5 mg/kg GW274150, and iNOS-/-) were subjected to bilateral renal ischemia (30 min) followed by reperfusion (24 h) after which renal dysfunction (serum urea, creatinine), renal myeloperoxidase (MPO) activity and malondialdehyde (MDA) levels were measured. GW274150, administered prior to I/R, significantly reduced serum urea, serum creatinine, AST, and NAG indicating reduction of renal dysfunction and injury caused by I/R. GW274150 reduced histol. evidence of tubular injury and markedly reduced immunohistochem. evidence of nitrotyrosine and PAR formation, indicating reduced peroxynitrite formation and poly (ADP-ribose) polymerase (PARP) activation, resp. GW274150 abolished the rise in the plasma levels of nitrate (indicating reduced NO production). GW274150 also reduced the renal dysfunction in wild-type mice to levels similar to that observed in iNOS-/- mice subjected to I/R. Renal MPO activity and MDA levels were significantly reduced in wild-type mice administered GW274150 and iNOS-/- mice subjected to renal I/R, indicating reduced polymorphonuclear leukocyte (PMN) infiltration and lipid peroxidn. These results suggest that (1) an enhanced formation of NO by iNOS contributes to the pathophysiol. of renal I/R injury and (2) GW274150 reduces I/R injury of the kidney. We propose that selective inhibitors of iNOS activity may be useful against renal dysfunction and injury associated with I/R of the kidney.
IT 210354-22-6, GW274150

RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(GW274150, selective inhibitor of iNOS, reduces exptl. renal ischemia/reperfusion injury)

RN 210354-22-6 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 67 THERE ARE 67 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 11 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2003:51504 CAPLUS

DN 139:159864

TI A novel, potent and selective inhibitor of the activity of inducible nitric oxide synthase (GW274150) reduces the organ injury in hemorrhagic shock

AU McDonald, M. C.; Izumi, M.; Cuzzocrea, S.; Thiemermann, C.

CS The William Harvey Research Institute, St. Bartholomew's and The Royal London School of Medicine and Dentistry, London, EC1M6BQ, UK

SO Journal of Physiology and Pharmacology (2002), 53(4, Pt. 1), 555-569
CODEN: JPHPEI; ISSN: 0867-5910

PB Polish Physiological Society

DT Journal

LA English

AB An enhanced formation of nitric oxide (NO) by the inducible NO synthase (iNOS) may contribute to the pathophysiol. of hemorrhagic shock. This study investigates the effect of a novel, potent and selective inhibitor of iNOS activity (GW274150) on the circulatory failure and the organ injury and dysfunction associated with hemorrhagic shock in the anesthetized rat. Hemorrhage (sufficient to lower mean arterial blood pressure to 45 mmHg for 90 min) and subsequent resuscitation with shed blood resulted (within 4 h after resuscitation) in a delayed fall in blood pressure, renal and liver injury and dysfunction as well as the pancreatic injury. Pre-treatment of rats with GW274150 (5 mg/kg at 30 min prior to the onset of hemorrhage) attenuated the renal dysfunction as well as the liver and pancreatic injury caused by hemorrhage and resuscitation. Interestingly, GW274150 did not reduce the delayed fall in blood pressure associated with hemorrhagic shock. We propose that an enhanced formation of NO from iNOS contributes to the organ injury and dysfunction in hemorrhagic shock, and that highly selective inhibitors of iNOS activity, such as GW274150, may represent a novel therapeutic approach for the therapy of hemorrhagic shock.

IT 210354-22-6, GW274150


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(novel, potent and selective inhibitor of activity of inducible nitric oxide synthase (GW274150) reduces the organ injury in hemorrhagic shock)

RN 210354-22-6 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 12 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2002:804285 CAPLUS

DN 138:314136

TI Beneficial effects of GW274150, a novel, potent and selective inhibitor of iNOS activity, in a rodent model of collagen-induced arthritis

AU Cuzzocrea, Salvatore; Chatterjee, Prabal K.; Mazzon, Emanuela; McDonald, Michelle C.; Dugo, Laura; Di Paola, Rosanna; Serraino, Ivana; Britti, Domenico; Caputi, Achille P.; Thiemermann, Christoph

CS School of Medicine, Institute of Pharmacology, University of Messina, Policlinico Universitario, Gazzi, Messina, 98100, Italy

SO European Journal of Pharmacology (2002), 453(1), 119-129

CODEN: EJPHAZ; ISSN: 0014-2999

PB Elsevier Science B.V.

DT Journal

LA English

AB The aim of this study was to investigate the role of inducible nitric oxide synthase (iNOS) on the modulation of the inflammatory response in mice subjected to collagen-induced arthritis. Collagen-induced arthritis was induced in wild-type mice (iNOS-WT) treated with GW274150, a novel, potent and selective inhibitor of iNOS activity, and in mice lacking the gene for iNOS (iNOS knock-out, iNOS-KO), by an intradermal injection of 100 µl of emulsion containing 100 µg of bovine type II collagen and complete Freund's adjuvant at the base of the tail. After 21 days, a second injection of type II collagen in complete Freund's adjuvant was administered. iNOS-WT mice developed erosive hind paw arthritis when immunized with type II collagen in complete Freund's adjuvant. Over a 35-day period, macroscopic clin. evidence of collagen-induced arthritis first appeared as periarticular erythema and edema in the hind paws. By day 28, the incidence of collagen-induced arthritis was 100% in type II collagen-challenged iNOS-WT mice and the severity of collagen-induced arthritis progressed with radiog. evaluation revealing resorption of bone. Histopathol. of collagen-induced arthritis mice demonstrated erosion of the cartilage at the joint margins. iNOS-WT mice treated with GW274150 (5 mg/kg, i.p. daily) starting at the onset of arthritis (day 23), and iNOS-KO mice showed a delay of the development of the clin. signs at days 24-35 and an improvement of the histol. status in the knee and paw. Immunohistochem. anal. for nitrotyrosine and for poly(ADP-ribose) polymerase revealed pos. staining in inflamed joints from type II collagen-treated iNOS-WT mice. The degree of staining for nitrotyrosine and poly(ADP-ribose) polymerase were markedly reduced in tissue sections obtained from type II collagen-treated iNOS-WT mice, who had received GW274150 and from iNOS-KO mice. Furthermore, radiog. signs of protection against bone resorption were present in the joints of iNOS-WT mice treated with GW274150 as well as in the joint from iNOS-KO mice. This study provides the first evidence that GW274150, a novel, potent and selective inhibitor of iNOS activity, attenuates the degree of chronic inflammation and tissue damage associated with collagen-induced arthritis in mice. Furthermore, these results suggest that the induction of iNOS and NO production are essential for the up-regulation of the inflammatory response during exptl. collagen-induced arthritis.

IT 210354-22-6, GW274150


RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(effects of GW274150 in a rodent model of collagen-induced arthritis)

RN 210354-22-6 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RE.CNT 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 13 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2001:597331 CAPLUS

DN 136:288829

TI GW274150 is a potent, long-acting, highly-selective inhibitor of inducible nitric oxide synthase (NOS-2) with therapeutic potential in post-operative ileus

AU Alderton, W.; Angell, A.; Clayton, N.; Craig, C.; Dawson, J.; Frend, A.; McGill, J.; Mangel, A.; Moncada, S.; Rees, D.; Russell, L.; Schwartz, S.; Waslidge, N.; Knowles, R.

CS Glaxo Wellcome R and D, Stevenage, SG1 2NY, UK

SO Portland Press Proceedings (2000), 16(Biology of Nitric Oxide, Part 7), 22
CODEN: POPPEF; ISSN: 0966-4068

PB Portland Press Ltd.

DT Journal

LA English

AB GW274150 [(S)-2-amino-7-acetamidino-5-thioheptanoic acid] is a novel α -amino acid that potently inhibited human inducible nitric oxide synthase (iNOS) with selectivity vs. human eNOS and nNOS. In studies with purified NOS isoforms, GW274150 was a time-dependent, arginine-site inhibitor of iNOS and a rapidly-reversible inhibitor of eNOS. This novel compound had a long pharmacokinetic half-life and high oral bioavailability in several species. The selectivity of GW274150 against the constitutive NOS isoforms was maintained in vivo, the compound producing no significant effect on conscious mouse blood pressure dosed at 100 mg/kg and on rat brain plus nitrite levels at 50 mg/kg. Post-operative ileus is one potential therapeutic application for GW274150. In a rat model of post-operative ileus, GW274150 was maximally effective at 1-5 mg/kg, yielding a 67% reversal of delayed GI transit. The compound was also effective in a rat model of acute inflammatory pain (adjuvant).

IT 210354-22-6, GW 274150


RL: PAC (Pharmacological activity); PKT (Pharmacokinetics); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(GW274150 is a potent, long-acting, highly-selective inhibitor of inducible nitric oxide synthase (NOS-2) with therapeutic potential in post-operative ileus)

RN 210354-22-6 CAPLUS

CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 14 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
AN 2000:209102 CAPLUS
DN 133:12344
TI Inhibition of inducible nitric oxide synthase by acetamidine derivatives
of hetero-substituted lysine and homolysine
AU Young, Robert J.; Beams, Richard M.; Carter, Keith; Clark, Helen A. R.;
Coe, Diane M.; Chambers, C. Lynn; Davies, P. Ifeyinwa; Dawson, John;
Drysdale, Martin J.; Franzman, Karl W.; French, Colin; Hodgson, Simon T.;
Hodson, Harold F.; Kleanthous, Savvas; Rider, Peter; Sanders, Daniela;
Sawyer, David A.; Scott, Keith J.; Shearer, Barry G.; Stocker, Richard;
Smith, Steven; Tackley, Miriam C.; Knowles, Richard G.
CS Glaxo Wellcome Research and Development, Stevenage, SG1 2NY, UK
SO Bioorganic & Medicinal Chemistry Letters (2000), 10(6), 597-600
CODEN: BMCLE8; ISSN: 0960-894X
PB Elsevier Science Ltd.
DT Journal
LA English
AB The synthesis and in vitro evaluation of the acetamidine derivs. of
hetero-substituted lysine and homolysine analogs have identified potent
inhibitors of human nitric oxide synthase enzymes, including examples with
marked selectivity for the inducible isoform.
IT 210354-22-6, GW 274150
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); PRP (Properties); BIOL (Biological study)
(inhibition of inducible nitric oxide synthase by acetamidine derivs.
of hetero-substituted lysine and homolysine)
RN 210354-22-6 CAPLUS
CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RE.CNT 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 15 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
AN 1999:753054 CAPLUS
DN 131:346497
TI Use of nitric oxide synthase inhibitors in the manufacture of a medicament
for the prophylaxis or treatment of bacterial infection
IN Alderton, Wendy Karen; Knowles, Richard Graham; Ladel, Christoph Hubertus
PA Glaxo Group Limited, UK
SO PCT Int. Appl., 34 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9959566	A1	19991125	WO 1999-EP3265	19990512
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ,				
DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS,				
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK,				
MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ,				
TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ,				

MD, RU, TJ, TM
 RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK,
 ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG,
 CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
 AU 9940406 A1 19991206 AU 1999-40406 19990512
 PRAI GB 1998-10299 A 19980515
 WO 1999-EP3265 W 19990512
 OS MARPAT 131:346497
 AB Inducible nitric oxide synthase inhibitors are used for the manufacture of a
 medicament for the prophylaxis or treatment of a bacterial infection,
 where the inhibitor of inducible nitric oxide synthase is e.g.
 HN:C(R1)NHR2 [R1 = C1-6 straight or branched chain alkyl; Q = QC(NH2)CO2H
 (Q = alkylene, alkenylene, etc.), ring-substituted benzyl] or a
 pharmaceutically acceptable salt, ester, or amide thereof.
 IT 210354-22-6
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES
 (Uses)
 (nitric oxide synthase inhibitors for prophylaxis or treatment of
 bacterial infection)
 RN 210354-22-6 CAPLUS
 CN L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L1 ANSWER 16 OF 16 CAPLUS COPYRIGHT 2006 ACS on STN
 AN 1998:490618 CAPLUS
 DN 129:122862
 TI Preparation of S-[2-(1-iminoethylamino)ethyl]homocysteine as nitric oxide
 synthase inhibitor
 IN Beams, Richard Mansfield; Drysdale, Martin James; Franzman, Karl Witold;
 Frend, Anthony Joseph; Hodson, Harold Francis; Knowles, Richard Graham;
 Rees, Daryl David; Sawyer, David Alan
 PA Glaxo Group Ltd., UK; Beams, Richard Mansfield; Drysdale, Martin James;
 Franzman, Karl Witold; Frend, Anthony Joseph; Hodson, Harold Francis;
 Knowles, Richard Graham; Rees, Daryl David; Sawyer, David Alan
 SO PCT Int. Appl., 29 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9830537	A1	19980716	WO 1998-EP96	19980109
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2277877	AA	19980716	CA 1998-2277877	19980109
AU 9862083	A1	19980803	AU 1998-62083	19980109

AU 723095	B2	20000817		
ZA 9800179	A	19990709	ZA 1998-179	19980109
EP 958277	A1	19991124	EP 1998-904050	19980109
EP 958277	B1	20011121		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
EE 9900281	A	20000215	EE 1999-281	19980109
EE 4013	B1	20030415		
JP 2000504041	T2	20000404	JP 1998-530549	19980109
JP 3251301	B2	20020128		
BR 9806870	A	20000418	BR 1998-6870	19980109
NZ 336379	A	20010126	NZ 1998-336379	19980109
AT 209183	E	20011215	AT 1998-904050	19980109
PT 958277	T	20020531	PT 1998-904050	19980109
ES 2168737	T3	20020616	ES 1998-904050	19980109
SK 283201	B6	20030304	SK 1999-933	19980109
AP 1204	A	20030915	AP 1999-1603	19980109
W: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW				
IL 130551	A1	20040104	IL 1998-130551	19980109
CZ 293099	B6	20040218	CZ 1999-2483	19980109
PL 189973	B1	20051031	PL 1998-334368	19980109
TW 502010	B	20020911	TW 1998-87100434	19980114
TW 538021	B	20030621	TW 1999-88103866	19980114
NO 9903429	A	19990712	NO 1999-3429	19990712
NO 312192	B1	20020408		
US 6369272	B1	20020409	US 1999-341220	19990824
HK 1021531	A1	20020315	HK 2000-100440	20000124
US 2002010366	A1	20020124	US 2001-930605	20010815
US 6620848	B2	20030916		
PRAI US 1997-69882P	P	19970113		
US 1997-783402	A	19970113		
WO 1998-EP96	W	19980109		
US 1999-341220	A1	19990824		
OS	MARPAT 129:122862			
AB	HN:CMcNHCH2CH2SCH2CH2CH(NH2)CO2H (I) was prepared for use as a selective inhibitor of nitric oxide synthase (NOS). Thus, (S)-I was prepared by treatment of L-homocystine with Na in liquid NH3 and then N-benzyloxycarbonylethanolamine tosylate, cleavage of the benzyloxycarbonyl protecting group with HBr in AcOH, and reaction with Et acetimidate hydrochloride. (S)-I was assayed for inhibition of inducible and endothelial NOS (IC50 = 0.73 and 43 µM, resp.).			
IT	210354-22-6P RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of [2-(1-iminoethylamino)ethyl]homocysteine as nitric oxide synthase inhibitor)			
RN	210354-22-6 CAPLUS			
CN	L-Homocysteine, S-[2-[(1-iminoethyl)amino]ethyl]- (9CI) (CA INDEX NAME)			

Absolute stereochemistry.



RE.CNT 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	92.10	92.31
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-12.00	-12.00

STN INTERNATIONAL LOGOFF AT 08:08:07 ON 19 SEP 2006